

Intersecting Families of Separated Sets

John Talbot
Merton College
University of Oxford
E-mail: talbot@maths.ox.ac.uk

March 10th 2002

Abstract

A set $A \subseteq \{1, 2, \dots, n\}$ is said to be *k-separated* if, when considered on the circle, any two elements of A are separated by a gap of size at least k .

We prove a conjecture due to Holroyd and Johnson [3],[4] that an analogue of the Erdős-Ko-Rado theorem holds for *k-separated* sets. In particular the result holds for the vertex-critical subgraph of the Kneser graph identified by Schrijver [7], the collection of separated sets. We also give a version of the Erdős-Ko-Rado theorem for weighted *k-separated* sets.

1 Introduction

A family of sets is *intersecting* if any two sets from the family meet. If $[n]^{(r)}$ is the collection of all r -sets from $[n] = \{1, 2, \dots, n\}$ how large can an intersecting family of sets $\mathcal{A} \subseteq [n]^{(r)}$ be? If $n < 2r$ this is easy to answer since $[n]^{(r)}$ is intersecting. However, for $n \geq 2r$ this question is more difficult. It was answered by Erdős, Ko and Rado [1].

Theorem 1 (Erdős, Ko and Rado [1]) *Let $n \geq 2r$ and $\mathcal{A} \subset [n]^{(r)}$ be intersecting. Then $|\mathcal{A}| \leq |\mathcal{A}_1|$, where $\mathcal{A}_1 = \{A \in [n]^{(r)} : 1 \in A\}$.*

Mathematics Subject Classification: 05D05 Extremal Set Theory, 05C65 Hypergraphs.

When $n \geq 2r + 1$ the above result can be extended to show that equality holds iff $\mathcal{A} \simeq \mathcal{A}_1$.

Many questions concerning families of sets from $[n]^{(r)}$ can be framed in the language of graphs. Consider the *Kneser graph*, $K_{n,r}$, with vertex set $[n]^{(r)}$ and edges between any two vertices corresponding to disjoint r -sets. The Erdős-Ko-Rado theorem (Theorem 1) can be restated as: the largest independent set of vertices in $K_{n,r}$ has order $\binom{n-1}{r-1}$.

One of the most fundamental properties of a graph is its chromatic number. A longstanding conjecture due to Kneser [5] was that the chromatic number of $K_{n,r}$ is $n - 2r + 2$. This was answered in the affirmative by Lovász in 1977 [6]. Later Schrijver [7] identified a vertex-critical subgraph of $K_{n,r}$, that is a minimal subgraph of $K_{n,r}$ with chromatic number $n - 2r + 2$. In order to describe this subgraph we require the following definition. We say that a set $A \in [n]^{(r)}$ is *separated* if, when considered as a subset of $[n]$ arranged around a circle in the usual ordering, A does not contain any two adjacent points. Schrijver's vertex-critical subgraph of the Kneser graph is the subgraph induced by those vertices corresponding to the collection of all separated sets in $[n]^{(r)}$.

Let us denote the collection of all separated sets in $[n]^{(r)}$ by $[n]_*^{(r)}$. Then the corresponding subgraph of the Kneser graph has (by Schrijver's result) chromatic number $n - 2r + 2$. However, the independence number of this subgraph was previously not known. This was a rather curious situation since generally determining the independence number of a graph is "easier" than determining its chromatic number.

For the remainder of this paper we will consider a well-known conjecture of Holroyd and Johnson on this problem, namely that an analogue of the Erdős-Ko-Rado theorem holds for intersecting families of separated sets. In fact their conjecture is more general. They define for any integer $k \geq 1$ the collection of k -separated r -sets in $[n]^{(r)}$ to be those r -sets $A = \{a_1, \dots, a_r\}$ with $a_1 < a_2 < \dots < a_r$ satisfying $a_{i+1} - a_i > k$ for $i = 1, \dots, r$, where $a_{r+1} = a_1 + n$. We will denote this family by $[n]_k^{(r)}$. Note that a 1-separated set is simply a separated set.

Conjecture 1 (Holroyd and Johnson [3],[4]) Let n , k and r be positive integers satisfying $n \geq (k + 1)r$. Suppose $\mathcal{A} \subseteq [n]_k^{(r)}$ is intersecting. Then $|\mathcal{A}| \leq |\mathcal{A}_1^*|$, where $\mathcal{A}_1^* = \{A \in [n]_k^{(r)} : 1 \in A\}$.

Our main result, Theorem 3, is a proof of this conjecture. The key idea used in the proof is a type of "compression". Standard proofs using compression

(such as the original proof of the Erdős-Ko-Rado theorem [1]) generally rely on certain properties of sets being preserved under this operation. Our proof is quite different since the property of being k -separated is not preserved under our compression operation.

In the next section of this paper we prove the conjecture for separated sets (the case $k=1$). This proof is then easily generalised in the subsequent section to yield the full result. We also characterise the extremal families.

In the final section of this paper we give a new version of the Erdős-Ko-Rado theorem for weighted k -separated sets.

2 The Erdős-Ko-Rado Theorem for separated sets

Theorem 2 *Let $n \geq 2r$ and $\mathcal{A} \subseteq [n]_*^{(r)}$ be intersecting then $|\mathcal{A}| \leq |\mathcal{A}_1^*|$ where $\mathcal{A}_1^* = \{A \in [n]_*^{(r)} : 1 \in A\}$. Moreover, for $n \neq 2r + 2$ the extremal family $\mathcal{A}_1^*$ is unique up to isomorphism.*

If $n = 2r + 2$ then other extremal families exist. For example if $d = \lfloor r/2 \rfloor$ then for each $1 \leq i \leq d$ the following family is extremal

$$\mathcal{B}_i = \{A \in [n]_*^{(r)} : |A \cap \{1, 3, \dots, 4i + 1\}| \geq i + 1\}.$$

Before giving the proof in full we sketch the basic ideas used.

The proof uses induction on n , where $\mathcal{A} \subseteq [n]_*^{(r)}$ is our intersecting family. The base case ($n = 2r$) is trivial so we may assume that $n \geq 2r + 1$. We show that $|\mathcal{A}| = |\mathcal{A}_1| + |\mathcal{A}_2|$, where $\mathcal{A}_1 \subseteq [n - 1]_*^{(r)}$ and $\mathcal{A}_2 \subseteq [n - 2]_*^{(r-1)}$ are both intersecting. Since $n \geq 2r + 1$ the inductive hypothesis applies in both cases and this yields the desired result.

In order to describe the families $\mathcal{A}_1$ and $\mathcal{A}_2$ we introduce a “compression” function, f , that maps points in $[n] \setminus \{1\}$ anticlockwise by one and leaves 1 fixed. Then $\mathcal{A}_1$ consists of those sets in $f(\mathcal{A})$ that are still separated (as subsets of $[n - 1]$). Clearly $\mathcal{A}_1$ is an intersecting family in $[n - 1]_*^{(r)}$.

If $A \in [n]_*^{(r)}$ but $f(A) \notin [n - 1]_*^{(r)}$ then A must either contain the pair $\{1, 3\}$ or the pair $\{2, n\}$. So let $\mathcal{D}$ consist of those sets in $\mathcal{A}$ containing $\{1, 3\}$ and let $\mathcal{E}$ consist of those sets in $\mathcal{A}$ containing $\{2, n\}$. Applying the compression to both of these families we see that each set in $f(\mathcal{D})$ contains $\{1, 2\}$ and each set in $f(\mathcal{E})$ contains $\{1, n - 1\}$. We now remove the point 1 from each of these

compressed sets to give a new family $\mathcal{G} = (f(\mathcal{D}) - \{1\}) \cup (f(\mathcal{E}) - \{1\})$. Clearly this is a disjoint union, the fact that $\mathcal{G}$ is also intersecting is less obvious.

We now need to consider those cases when two distinct sets $A, B \in \mathcal{A}$ both compress to the same set (i.e. $f(A) = f(B)$). It is easy to see that this can only happen if $A \Delta B = \{1, 2\}$. Hence $|\mathcal{D}| = |f(\mathcal{D})|$ and $|\mathcal{E}| = |f(\mathcal{E})|$. Also any set in $\mathcal{A}_1$ has at most two preimages in $\mathcal{A}$. Let $\mathcal{F}$ consist of those sets in $\mathcal{A}_1$ that have two preimages in $\mathcal{A}$ then each set in $\mathcal{F}$ contains 1. Consider the new family $\mathcal{H} = (\mathcal{F} - \{1\}) \cup \mathcal{G}$. Then $\mathcal{H}$ is the union of disjoint families (to see this just check that if $H \in \mathcal{H}$ then $H \cap \{2, n-1\} = \emptyset \iff H \in \mathcal{F} - \{1\}$). Furthermore $\mathcal{H}$ is also intersecting. Now no set in $\mathcal{H}$ contains 1 and so $|\mathcal{H}| = |f(\mathcal{H})|$. Applying our compression function to $\mathcal{H}$ we define $\mathcal{A}_2 = f(\mathcal{H})$. We then find that $\mathcal{A}_2$ is in fact an intersecting family in $[n-2]_*^{(r-1)}$.

Finally $|\mathcal{A}| = |\mathcal{A}_1| + |\mathcal{F}| + |\mathcal{D}| + |\mathcal{E}| = |\mathcal{A}_1| + |\mathcal{F}| + |\mathcal{G}| = |\mathcal{A}_1| + |\mathcal{H}| = |\mathcal{A}_1| + |\mathcal{A}_2|$ as claimed.

In order to show the uniqueness of the extremal family (for $n \neq 2r+2$) we note first that the result holds trivially for $n = 2r$ and $n = 2r+1$ so we may assume that $n \geq 2r+3$. We then proceed by induction on r . Clearly the result holds for $r = 1$ so we proceed to the inductive step. Consider the family $\mathcal{A}_2$ defined above. For equality to hold we must have $|\mathcal{A}_2| = \binom{n-r-2}{r-2}$ and since $n-2 \geq 2(r-1)+3$ our inductive hypothesis for $r-1$ implies that there exists $i \in [n]$ such that every set in $\mathcal{A}_2$ contains i . Considering the different possibilities for i we find that either $\mathcal{A} \simeq \mathcal{A}_1^*$ or there exists $j \in [n]$ such that $\{A \in [n]_*^{(r)} : j, j+2 \in A\} \subset \mathcal{A}$. Without loss of generality we may suppose that $j = 1$. This easily implies that every set in $\mathcal{A}$ must contain either 1 or 3. Partition $\mathcal{A}$ as $\mathcal{A} = \mathcal{B}_1 \cup \mathcal{B}_3 \cup \mathcal{B}_{1,3}$, where $\mathcal{B}_1 = \{A \in \mathcal{A} : 1 \in A, 3 \notin A\}$, $\mathcal{B}_3 = \{A \in \mathcal{A} : 1 \notin A, 3 \in A\}$ and $\mathcal{B}_{1,3} = \{A \in \mathcal{A} : 1, 3 \in A\}$. To prove that $\mathcal{A} \simeq \mathcal{A}_1^*$ it is sufficient to show that one of the families $\mathcal{B}_1$ or $\mathcal{B}_3$ must be empty. It can be shown that if $n \geq 2r+3$ and both families are non-empty then $\mathcal{A}$ contains two disjoint sets, a contradiction.

Proof of Theorem 2. We proceed by induction on n . First note that the result is trivial for $n \leq 4$ so we may suppose $n \geq 5$. Also we may suppose that $n \geq 2r+1$ since if $n = 2r$ then there are only two separated sets and these are disjoint so the result follows.

Let $\mathcal{A} \subseteq [n]_*^{(r)}$ be intersecting. Consider the following partition of $\mathcal{A}$

$$\mathcal{A} = \mathcal{B} \cup \mathcal{C} \cup \mathcal{D} \cup \mathcal{E},$$

where

$$\begin{aligned}\mathcal{B} &= \{A \in \mathcal{A} : 1 \notin A \text{ and } (2 \notin A \text{ or } n \notin A)\}, \\ \mathcal{C} &= \{A \in \mathcal{A} : 1 \in A \text{ and } 3 \notin A\}, \\ \mathcal{D} &= \{A \in \mathcal{A} : 1, 3 \in A\}, \\ \mathcal{E} &= \{A \in \mathcal{A} : 2, n \in A\}.\end{aligned}$$

Define the function $f : [n] \rightarrow [n-1]$ by

$$f(j) = \begin{cases} 1, & j = 1 \\ j-1, & j \geq 2. \end{cases}$$

We will need to use the following results (their proofs are given later).

Lemma 1 *If $\mathcal{A}_1^* = \{A \in [n]_*^{(r)} : 1 \in A\}$ then*

$$|\mathcal{A}_1^*| = \binom{n-r-1}{r-1}.$$

Lemma 2 *If $A, B \in [n]_*^{(r)}$, $A \neq B$ and f is as defined above then*

$$f(A) = f(B) \implies A \Delta B = \{1, 2\}.$$

Lemma 3 *If $A \in \mathcal{B} \cup \mathcal{C}$ and $\mathcal{B}, \mathcal{C}, f$ are as defined above then $f(A) \in [n-1]_*^{(r)}$.*

Lemma 4 *If $\mathcal{I}$ is an intersecting family of sets then so is $f(\mathcal{I})$.*

Since no set in $\mathcal{B}$ contains 1 and no set in $\mathcal{C}$ contains 2, Lemma 2 implies that $|f(\mathcal{B})| = |\mathcal{B}|$ and $|f(\mathcal{C})| = |\mathcal{C}|$. Hence

$$|\mathcal{B}| + |\mathcal{C}| = |f(\mathcal{B})| + |f(\mathcal{C})| = |f(\mathcal{B}) \cup f(\mathcal{C})| + |f(\mathcal{B}) \cap f(\mathcal{C})|.$$

Then, since $\mathcal{B} \cup \mathcal{C}$ is an intersecting family in $[n]_*^{(r)}$, Lemmas 3 and 4 imply that $f(\mathcal{B}) \cup f(\mathcal{C}) = f(\mathcal{B} \cup \mathcal{C})$ is an intersecting family in $[n-1]_*^{(r)}$. Hence by our inductive hypothesis for $n-1 \geq 2r$ and Lemma 1 we have

$$|f(\mathcal{B}) \cup f(\mathcal{C})| \leq \binom{n-1-r-1}{r-1} = \binom{n-r-2}{r-1}.$$

Let

$$\mathcal{F} = f(\mathcal{B}) \cap f(\mathcal{C}).$$

For any family of sets $\mathcal{G}$ let

$$\mathcal{G} - \{1\} = \{G \setminus \{1\} : G \in \mathcal{G}\}.$$

Define

$$\mathcal{H} = (f(\mathcal{D}) - \{1\}) \cup (f(\mathcal{E}) - \{1\}) \cup (\mathcal{F} - \{1\}).$$

We now require the following result.

Lemma 5 *Let $\mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{H}$ and f be as defined above, then*

- (a) $\mathcal{H} \subseteq [n-1]_*^{(r-1)}$.
- (b) $f(\mathcal{D}) - \{1\}$, $f(\mathcal{E}) - \{1\}$ and $\mathcal{F} - \{1\}$ are pairwise disjoint families of sets.
- (c) $\mathcal{H}$ is intersecting.
- (d) $f(\mathcal{H}) \subseteq [n-2]_*^{(r-1)}$.

Since no set in $\mathcal{H}$ contains 1 Lemma 2 implies that $|\mathcal{H}| = |f(\mathcal{H})|$. Lemma 5(c) says that $\mathcal{H}$ is intersecting and so Lemma 4 implies that $f(\mathcal{H})$ is also intersecting. Moreover Lemma 5(d) tells us that $f(\mathcal{H}) \subseteq [n-2]_*^{(r-1)}$. So using Lemma 1 and the inductive hypothesis for $n-2 \geq 2(r-1)$ we obtain

$$|\mathcal{H}| \leq \binom{n-2-(r-1)-1}{(r-1)-1} = \binom{n-r-2}{r-2}.$$

Using Lemma 2 again we also have $|\mathcal{D}| = |f(\mathcal{D})|$ and $|\mathcal{E}| = |f(\mathcal{E})|$. Finally Lemma 5(b) tells us that

$$\begin{aligned} |\mathcal{H}| &= |f(\mathcal{D})| + |f(\mathcal{E})| + |\mathcal{F}| \\ &= |\mathcal{D}| + |\mathcal{E}| + |\mathcal{F}|. \end{aligned}$$

Hence

$$\begin{aligned} |\mathcal{A}| &= |\mathcal{B}| + |\mathcal{C}| + |\mathcal{D}| + |\mathcal{E}| \\ &= |f(\mathcal{B}) \cup f(\mathcal{C})| + |\mathcal{F}| + |\mathcal{D}| + |\mathcal{E}| \\ &= |f(\mathcal{B}) \cup f(\mathcal{C})| + |\mathcal{H}| \\ &\leq \binom{n-r-2}{r-1} + \binom{n-r-2}{r-2} = \binom{n-r-1}{r-1} = |\mathcal{A}_1^*|. \end{aligned} \tag{1}$$

This completes the proof of the bound in Theorem 2.

In order to prove that the extremal family is unique for $n \neq 2r+2$ we will require the following results (again their proofs follow later).

Lemma 6 *If equality holds in (1) and the family $f(\mathcal{H})$ defined above is isomorphic to $\mathcal{B}_1^* = \{A \in [n-2]_*^{(r-1)} : 1 \in A\}$ then either $\mathcal{A} \simeq \mathcal{A}_1^*$ or there exists $j \in [n]$ such that $\mathcal{A}_{j,j+2}^* = \{A \in [n]_*^{(r)} : j, j+2 \in A\} \subset \mathcal{A}$.*

Lemma 7 *If equality holds in (1) and $\mathcal{A}_{1,3}^* = \{A \in [n]_*^{(r)} : 1, 3 \in A\} \subset \mathcal{A}$ then every set in $\mathcal{A}$ meets $\{1, 3\}$. Moreover if $A = \{1, a_2, \dots, a_r\} \in [n]_*^{(r)}$ satisfies $A \cap \{1, 3\} = \{1\}$ then exactly one of the sets A and $g(A) = \{3, a_2+1, \dots, a_r+1\}$ belongs to $\mathcal{A}$.*

Suppose equality holds in (1). If $n = 2r$ or $n = 2r + 1$ it is easy to deduce that $\mathcal{A} \simeq \mathcal{A}_1^*$ so we may suppose that $n \geq 2r + 3$. We will prove that $\mathcal{A} \simeq \mathcal{A}_1^*$ by induction on r . Clearly the result holds for $r = 1$ or 2 so suppose $r \geq 3$. Since equality holds in (1) we must have $|f(\mathcal{H})| = \binom{n-r-2}{r-2}$ and so our inductive hypothesis for r implies that $f(\mathcal{H}) \simeq \mathcal{B}_1^* = \{A \in [n-2]_*^{(r-1)} : 1 \in A\}$. If $\mathcal{A} \not\simeq \mathcal{A}_1^*$ then Lemma 6 implies that there exists $j \in [n]$ such that $\mathcal{A}_{j,j+2}^* = \{A \in [n]_*^{(r)} : j, j+2 \in A\} \subset \mathcal{A}$. Without loss of generality we may suppose that $j = 1$.

By the first part of Lemma 7 we know that every set in $\mathcal{A}$ meets $\{1, 3\}$. Hence we may write $\mathcal{A} = \mathcal{B}_1 \cup \mathcal{B}_3 \cup \mathcal{B}_{1,3}$, where $\mathcal{B}_1 = \{A \in \mathcal{A} : 1 \in A, 3 \notin A\}$, $\mathcal{B}_3 = \{A \in \mathcal{A} : 3 \in A, 1 \notin A\}$ and $\mathcal{B}_{1,3} = \{A \in \mathcal{A} : 1, 3 \in A\}$. For uniqueness to hold we need to show that one of the families $\mathcal{B}_1$ or $\mathcal{B}_3$ must be empty. We will prove that if both families are non-empty then $\mathcal{A}$ contains two disjoint sets, a contradiction.

Since we are supposing that equality holds in (1), Lemma 7 tells us that if $A = \{1, a_2, \dots, a_r\} \in [n]_*^{(r)}$ and $A \cap \{1, 3\} = \{1\}$ then exactly one of the sets A and $g(A) = \{3, a_2+1, \dots, a_r+1\}$ belong to $\mathcal{A}$. We now wish to show that if $A \in \mathcal{B}_1$ and $B \in [n]_*^{(r)}$ is obtained from A by shifting one vertex of $A \setminus \{1\}$ one place clockwise then $B \in \mathcal{B}_1$. Suppose not, then by Lemma 7 we have $g(B) \in \mathcal{B}_3$. But $g(B) \cap A = \emptyset$, contradicting the fact that $\mathcal{A}$ is intersecting. Similarly if $A \in \mathcal{B}_3$ and $B \in [n]_*^{(r)}$ is obtained from A by shifting one vertex of $A \setminus \{3\}$ one place anti-clockwise then $B \in \mathcal{B}_3$.

Now suppose, for a contradiction, that both $\mathcal{B}_1$ and $\mathcal{B}_3$ are non-empty then by shifting points one by one we may suppose that $A = \{3, 5, \dots, 2r+1\} \in \mathcal{B}_3$ and $B = \{1, n-2r+3, \dots, n-3, n-1\} \in \mathcal{B}_1$. If n is odd these are disjoint and we have a contradiction. If n is even then either there is a set in $\mathcal{B}_1$ not containing $n-1$ and so $C = \{1, n-2r+2, \dots, n-4, n-2\} \in \mathcal{B}_1$ and $A \cap C = \emptyset$ or, assuming $2r+1 \leq n-2$, $D = \{1, 5, 7, \dots, 2r+1\} \notin \mathcal{A}$. If the latter holds then Lemma 7 implies that $g(D) = \{3, 6, 8, \dots, 2r+2\} \in \mathcal{B}_3$. But $B \cap g(D) = \emptyset$. This contradiction completes the proof of Theorem 2. $\square$

We now prove the lemmas. The definitions of $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{H}$ and f are as in the proof of Theorem 2.

Proof of Lemma 1. Let $A \in \mathcal{A}_1^* = \{A \in [n]_*^{(r)} : 1 \in A\}$, then A is described uniquely by the gaps $a_1, a_2, \dots, a_r \geq 1$ such that

$$A = \{1, 2 + a_1, 3 + a_1 + a_2, \dots, r + a_1 + \dots + a_{r-1}\},$$

and $\sum_{i=1}^r a_i = n - r$. So $|\mathcal{A}_1^*|$ is equal to the number of ways of choosing integers $b_1, b_2, \dots, b_r \geq 0$ with $\sum_{i=1}^r b_i = n - 2r$ which is simply

$$\binom{n - 2r + r - 1}{r - 1} = \binom{n - r - 1}{r - 1}.$$

□

Proof of Lemma 2. If $j \geq 2$ then $j \in f(A) \iff j+1 \in A$. Hence $f(A) = f(B)$ implies that

$$A \cap \{3, 4, \dots, n\} = B \cap \{3, 4, \dots, n\}.$$

So if $A, B \in [n]_*^{(r)}$ and $f(A) = f(B)$ but $A \neq B$ then it must be the case that one of them contains 1 and the other contains 2. □

Proof of Lemma 3. Clearly $f(A) \in [n-1]^{(r)}$ (since nothing maps to n). So if $f(A) \notin [n-1]_*^{(r)}$ then we must have either $1, 2 \in f(A)$ or $1, n-1 \in f(A)$. Hence either $1, 3 \in A$ or $2, n \in A$ and so $A \notin \mathcal{B} \cup \mathcal{C}$. □

Proof of Lemma 4. If $\mathcal{I}$ is an intersecting family and $A, B \in f(\mathcal{I})$ then there exist $C, D \in \mathcal{I}$ such that $A = f(C)$ and $B = f(D)$. Then $\emptyset \neq f(C \cap D) \subseteq f(C) \cap f(D) = A \cap B$. So $f(\mathcal{I})$ is intersecting. □

Proof of Lemma 5(a). Each set in $\mathcal{H}$ comes from applying f to a set in $[n]_*^{(r)}$. The only way such a set can fail to belong to $[n-1]_*^{(r)}$ is if it contains 1 but no set in $\mathcal{H}$ contains 1. Hence $\mathcal{H} \subseteq [n-1]_*^{(r)}$. □

Proof of Lemma 5(b). The fact that the three families of sets $f(\mathcal{D}) - \{1\}$, $f(\mathcal{E}) - \{1\}$ and $\mathcal{F} - \{1\}$ are pairwise disjoint follows from considering how sets in these families meet the set $\{2, n-1\}$.

If $A \in f(\mathcal{D}) - \{1\}$ then $2 \in A$ and $n-1 \notin A$. If $A \in f(\mathcal{E}) - \{1\}$ then $2 \notin A$ and $n-1 \in A$. If $A \in \mathcal{F} - \{1\}$ then there exist $B \in \mathcal{B}$ and $C \in \mathcal{C}$ such that $f(B) = f(C) = A \cup \{1\}$. Then using Lemma 2 we have $C = (B \setminus \{2\}) \cup \{1\}$. So

$n, 3 \notin B$, which implies that $2, n-1 \notin A$. Hence the three families are pairwise disjoint since for any $H \in \mathcal{H}$ we have

$$H \cap \{2, n-1\} = \begin{cases} \{2\}, & H \in f(\mathcal{D}) - \{1\} \\ \{n-1\}, & H \in f(\mathcal{E}) - \{1\} \\ \emptyset, & H \in \mathcal{F} - \{1\}. \end{cases}$$

□

Proof of Lemma 5(c). We wish to show that

$$\mathcal{H} = (f(\mathcal{D}) - \{1\}) \cup (f(\mathcal{E}) - \{1\}) \cup (\mathcal{F} - \{1\}),$$

is intersecting.

Let $A, B \in \mathcal{H}$, there are six cases to examine. We consider first the three cases when A and B belong to the same subfamily of $\mathcal{H}$. If $A, B \in f(\mathcal{D}) - \{1\}$ then $2 \in A \cap B$. Similarly if $A, B \in f(\mathcal{E}) - \{1\}$ then $n-1 \in A \cap B$.

If $A, B \in \mathcal{F} - \{1\}$ then there are $B_1 \in \mathcal{B}$ and $C_1 \in \mathcal{C}$ such that $A = f(C_1) \setminus \{1\}$ and $B = f(B_1) \setminus \{1\}$. Then $C_1, B_1 \in \mathcal{A} \implies C_1 \cap B_1 \neq \emptyset$. Also $1 \notin B_1$ and $2 \notin C_1$ so there exists $j \in C_1 \cap B_1$ with $j \geq 3$. Hence $f(j) \in (f(C_1) \setminus \{1\}) \cap (f(B_1) \setminus \{1\}) = A \cap B$.

Now suppose $A \in f(\mathcal{D}) - \{1\}$ and $B \in f(\mathcal{E}) - \{1\}$. Then there exist $D \in \mathcal{D}$ and $E \in \mathcal{E}$ such that $A = f(D) \setminus \{1\}$ and $B = f(E) \setminus \{1\}$. Now $D, E \in \mathcal{A} \implies D \cap E \neq \emptyset$. Also $2 \notin D$ and $1 \notin E$ so there exists $j \in D \cap E$ with $j \geq 3$. Hence $f(j) \in A \cap B$.

Next suppose $A \in f(\mathcal{D}) - \{1\}$ and $B \in \mathcal{F} - \{1\}$. Then there exist $D \in \mathcal{D}$, $B_1 \in \mathcal{B}$ such that $A = f(D) \setminus \{1\}$ and $B = f(B_1) \setminus \{1\}$. Then $B_1, D \in \mathcal{A} \implies B_1 \cap D \neq \emptyset$. Also $1 \notin B_1$ and $2 \notin D$ so there exists $j \in B_1 \cap D$ with $j \geq 3$. Hence $f(j) \in A \cap B$.

Finally if $A \in f(\mathcal{E}) - \{1\}$ and $B \in \mathcal{F} - \{1\}$, then there exist $E \in \mathcal{E}$ and $C_1 \in \mathcal{C}$ such that $A = f(E) \setminus \{1\}$ and $B = f(C_1) \setminus \{1\}$. Since $C_1, E \in \mathcal{A}$ we have $C_1 \cap E \neq \emptyset$. Also $2 \notin C_1$ and $1 \notin E$ so there exists $j \in C_1 \cap E$ with $j \geq 3$. Then $f(j) \in A \cap B$.

Hence $\mathcal{H}$ is an intersecting family. □

Proof of Lemma 5(d). We need to prove that $f(\mathcal{H}) \subseteq [n-2]_*^{(r-1)}$. We note first that by part (a) of this lemma we have $\mathcal{H} \subseteq [n-1]_*^{(r-1)}$. Let $H \in \mathcal{H}$ and consider $f(H)$. Clearly $f(H) \in [n-2]^{(r-1)}$. We simply need to check that we do not have $1, 2 \in f(H)$ or $1, n-2 \in f(H)$. Now since $1 \notin H$ we have

$1 \in f(H) \iff 2 \in H$. But we showed during the proof of part (b) of this lemma that $2 \in H \implies H \in f(\mathcal{D}) - \{1\}$. In which case there is $D \in \mathcal{D}$ such that $H = f(D)$. Now $D \in \mathcal{D} \implies 1, 3 \in D \implies n, 4 \notin D \implies n-1, 3 \notin f(D) = H \implies 2, n-2 \notin f(H)$ as required. $\square$

Proof of Lemma 6. Suppose that $f(\mathcal{H}) \simeq \mathcal{B}_1^* = \{A \in [n-2]_*^{(r-1)} : 1 \in A\}$. We need to show that either $\mathcal{A} \simeq \mathcal{A}_1^*$ or there exists $j \in [n]$ such that $\mathcal{A}_{j,j+2}^* = \{A \in [n]_*^{(r)} : j, j+2 \in A\} \subset \mathcal{A}$.

Since $f(\mathcal{H}) \simeq \mathcal{B}_1^*$ there exists $i \in [n-2]$ such that $f(\mathcal{H}) = \mathcal{B}_i^* = \{A \in [n-2]_*^{(r-1)} : i \in A\}$. If $i = 1$ then every set in $\mathcal{H}$ contains 2 and so $\mathcal{H} = f(\mathcal{D}) - \{1\}$. Since $\mathcal{D} \subseteq \mathcal{A}_{1,3}^*$ and this last family has size $\binom{n-r-2}{r-2} = |f(\mathcal{H})| = |\mathcal{D}|$ then $\mathcal{D} = \mathcal{A}_{1,3}^*$. Similarly if $i = n-2$ then $\mathcal{E} = \mathcal{A}_{2,n}^*$. So if $i \in \{1, n-2\}$ then there is j such that $\mathcal{A}_{j,j+2}^* \subset \mathcal{A}$.

We will now show that if $i \in \{2, \dots, n-3\}$ then $\mathcal{A}_{1,i+2}^* \cup \mathcal{A}_{2,i+2}^* \subseteq \mathcal{A}$. Suppose $A = \{1, a_2, \dots, a_l, i+2, a_{l+2}, \dots, a_r\} \in \mathcal{A}_{1,i+2}^*$ but $A \notin \mathcal{A}$. Then since $f(\mathcal{H}) = \mathcal{B}_i^*$ it must be the case that $B = \{a_2-2, \dots, a_l-2, i, a_{l+2}-2, \dots, a_r-2\} \in f(\mathcal{H})$. Hence $C = (A \setminus \{1\}) \cup \{2\} \in \mathcal{A}$. Now $1 \in A$ implies that $n \notin C$ and so $C \notin \mathcal{D} \cup \mathcal{E}$. This implies that $B \in f(\mathcal{F})$ and so $A \in \mathcal{A}$, a contradiction. Hence $\mathcal{A}_{1,i+2}^* \subset \mathcal{A}$. Similarly $\mathcal{A}_{2,i+2}^* \subset \mathcal{A}$.

We can now show that every set in $\mathcal{A}$ must contain $i+2$ and so $\mathcal{A} \simeq \mathcal{A}_1^*$. Suppose not, then there exists a set $A = \{a_1, \dots, a_r\} \in \mathcal{A}$ such that $i+2 \notin A$. But then choosing a single point from each gap of A we may construct a set in $\mathcal{A}_{1,i+2}^* \cup \mathcal{A}_{2,i+2}^* \subseteq \mathcal{A}$, that is disjoint from A , contradicting the fact that $\mathcal{A}$ is intersecting. $\square$

Proof of Lemma 7. Suppose equality holds in (1) and $\mathcal{A}_{1,3}^* \subseteq \mathcal{A}$. We show first that every set in $\mathcal{A}$ meets $\{1, 3\}$. Suppose not, then there is a set $A = \{a_1, \dots, a_r\} \in \mathcal{A}$ such that $A \cap \{1, 3\} = \emptyset$. Then $a_3 \geq 6$ so $B = \{1, 3, a_3-1, \dots, a_r-1\} \in \mathcal{A}_{1,3}^* \subset \mathcal{A}$. But $A \cap B = \emptyset$ contradicting the fact that $\mathcal{A}$ is intersecting. Hence every set in $\mathcal{A}$ meets $\{1, 3\}$.

Let $\mathcal{B}_1 = \{A \in \mathcal{A} : 1 \in A, 3 \notin A\}$ and $\mathcal{B}_3 = \{A \in \mathcal{A} : 1 \notin A, 3 \in A\}$. For equality to hold in (1) we must have $|\mathcal{B}_1 \cup \mathcal{B}_3| = \binom{n-r-2}{r-1}$. Let $\mathcal{C}_1 = \{C \in [n]_*^{(r)} : 1 \in C, 3 \notin C\}$ and $\mathcal{C}_3 = \{C \in [n]_*^{(r)} : 1 \notin C, 3 \in C\}$. For $A = \{1, a_2, \dots, a_r\} \in \mathcal{C}_1$ define $g(A) = \{3, a_2+1, \dots, a_r+1\}$. Clearly $A \cap g(A) = \emptyset$ and g is a bijection from $\mathcal{C}_1$ to $\mathcal{C}_3$. Then, since $|\mathcal{C}_1| = |\mathcal{C}_3| = \binom{n-r-2}{r-1}$, it must be the case that for any $A \in \mathcal{C}_1$ exactly one of the sets A and $g(A)$ belongs to $\mathcal{A}$. $\square$

3 The Erdős-Ko-Rado Theorem for k -separated sets

Theorem 3 *Let $k \geq 2$, $r \geq 1$, $n \geq (k+1)r$ and $\mathcal{A} \subseteq [n]_k^{(r)}$ be intersecting then $|\mathcal{A}| \leq |\mathcal{A}_1^*|$ where $\mathcal{A}_1^* = \{A \in [n]_k^{(r)} : 1 \in A\}$. Moreover $\mathcal{A}_1^*$ is the unique extremal family up to isomorphism.*

Proof. We proceed by induction on n . First note that the result is trivial for $r \leq 2$ so we may suppose $r \geq 3$. Also we may suppose that $n \geq (k+1)r + 1$ since if $n = (k+1)r$ then there are only $k+1$ sets in $[n]_k^{(r)}$ and these are all pairwise disjoint so the result follows.

Let $\mathcal{A} \subseteq [n]_*^{(r)}$ be intersecting. Define the function $f : [n] \rightarrow [n-1]$ by

$$f(j) = \begin{cases} 1, & j = 1 \\ j-1, & j \geq 2. \end{cases}$$

Consider the following partition of $\mathcal{A}$

$$\mathcal{A} = \mathcal{B} \cup \mathcal{C} \cup \bigcup_{i=0}^k \mathcal{D}_i,$$

where

$$\begin{aligned} \mathcal{B} &= \{A \in \mathcal{A} : 1 \notin A \text{ and } f(A) \in [n-1]_k^{(r)}\}, \\ \mathcal{C} &= \{A \in \mathcal{A} : 1 \in A \text{ and } f(A) \in [n-1]_k^{(r)}\}, \\ \mathcal{D}_0 &= \{A \in \mathcal{A} : 1, k+2 \in A\}, \\ \mathcal{D}_i &= \{A \in \mathcal{A} : n+1-i, k+2-i \in A\}, \quad (1 \leq i \leq k). \end{aligned}$$

Note that this is a partition of $\mathcal{A}$ since either $f(A) \in [n-1]_k^{(r)}$ and so $A \in \mathcal{B} \cup \mathcal{C}$ or $f(A) \notin [n-1]_k^{(r)}$. If the latter holds then 1 must lie in a gap of A of size exactly k or 1 must be the left endpoint of such a gap and so there exists $0 \leq i \leq k$ such that $A \in \mathcal{D}_i$.

We will need to use the following results (again we defer their proofs until later).

Lemma 8 *If $\mathcal{A}_1^* = \{A \in [n]_k^{(r)} : 1 \in A\}$ then*

$$|\mathcal{A}_1^*| = \binom{n - kr - 1}{r - 1}.$$

Lemma 9 Suppose $A, B \in [n]_k^{(r)}$ with $A \neq B$ and f is as defined above. If $1 \leq j \leq k$ and f^j denotes f iterated j times then

$$f^j(A) = f^j(B) \implies A \Delta B = \{c, d\} \text{ for } 1 \leq c < d \leq j + 1.$$

Since no set in $\mathcal{B}$ contains 1, Lemma 9 with $j = 1$ implies that $|f(\mathcal{B})| = |\mathcal{B}|$. Similarly since no set in $\mathcal{C}$ contains 2 we have $|f(\mathcal{C})| = |\mathcal{C}|$. Hence

$$|\mathcal{B}| + |\mathcal{C}| = |f(\mathcal{B})| + |f(\mathcal{C})| = |f(\mathcal{B}) \cup f(\mathcal{C})| + |f(\mathcal{B}) \cap f(\mathcal{C})|.$$

Then, since $\mathcal{B} \cup \mathcal{C}$ is an intersecting family in $[n]_k^{(r)}$, Lemma 4 implies that $f(\mathcal{B}) \cup f(\mathcal{C}) = f(\mathcal{B} \cup \mathcal{C})$ is an intersecting family in $[n - 1]_k^{(r)}$. Hence by our inductive hypothesis for $n - 1 \geq (k + 1)r$ and Lemma 8 we have

$$|f(\mathcal{B}) \cup f(\mathcal{C})| \leq \binom{n - 1 - kr - 1}{r - 1} = \binom{n - kr - 2}{r - 1}.$$

Let

$$\mathcal{E} = f(\mathcal{B}) \cap f(\mathcal{C}).$$

For any family of sets $\mathcal{G}$ recall that

$$\mathcal{G} - \{1\} = \{G \setminus \{1\} : G \in \mathcal{G}\}.$$

Define

$$\mathcal{F} = (f^{k-1}(\mathcal{E}) - \{1\}) \cup \bigcup_{i=0}^k (f^k(\mathcal{D}_i) - \{1\}),$$

where f^j denotes the function f iterated j times.

We now require the following result.

Lemma 10 Let $\mathcal{D}_i, \mathcal{E}, \mathcal{F}$ and f be as defined above, then

- (a) $\mathcal{F} \subseteq [n - k]_k^{(r-1)}$.
- (b) $f^k(\mathcal{D}_0) - \{1\}, f^k(\mathcal{D}_1) - \{1\}, \dots, f^k(\mathcal{D}_k) - \{1\}$ and $f^{k-1}(\mathcal{E}) - \{1\}$ are pairwise disjoint families of sets.
- (c) $\mathcal{F}$ is intersecting.
- (d) $f(\mathcal{F}) \subseteq [n - k - 1]_k^{(r-1)}$.

Since no set in $\mathcal{F}$ contains 1, Lemma 9 with $j = 1$ implies that $|\mathcal{F}| = |f(\mathcal{F})|$. Lemma 10(c) says that $\mathcal{F}$ is intersecting and so Lemma 4 implies that $f(\mathcal{F})$ is also intersecting. Then Lemma 10(d) tells us that $f(\mathcal{F}) \subseteq [n - k - 1]_k^{(r-1)}$. So using Lemma 8 and the inductive hypothesis for $n - k - 1 \geq (k + 1)(r - 1)$ we obtain

$$|\mathcal{F}| \leq \binom{n - k - 1 - k(r - 1) - 1}{(r - 1) - 1} = \binom{n - kr - 2}{r - 2}.$$

If $D \in \mathcal{D}_i$ then the definition of $\mathcal{D}_i$ implies that

$$D \cap \{1, 2, \dots, k + 1\} = \begin{cases} \{1\}, & i = 0 \\ \{k + 2 - i\}, & i \geq 1. \end{cases}$$

Hence using Lemma 9 with $j = k$ we obtain $|\mathcal{D}_i| = |f^k(\mathcal{D}_i)|$ for $0 \leq i \leq k$.

If $E \in \mathcal{E} \subseteq f(\mathcal{C}) \subseteq [n - 1]_{k-1}^{(r)}$ then $E \cap \{1, 2, \dots, k\} = \{1\}$. So using Lemma 9 with $j = k - 1$ we obtain $|\mathcal{E}| = |f^{k-1}(\mathcal{E})|$. Finally Lemma 10(b) tells us that

$$\begin{aligned} |\mathcal{F}| &= |f^{k-1}(\mathcal{E})| + \sum_{i=0}^k |f^k(\mathcal{D}_i)| \\ &= |\mathcal{E}| + \sum_{i=0}^k |\mathcal{D}_i|. \end{aligned}$$

So we obtain

$$\begin{aligned} |\mathcal{A}| &= |\mathcal{B}| + |\mathcal{C}| + \sum_{i=0}^k |\mathcal{D}_i| \\ &= |f(\mathcal{B}) \cup f(\mathcal{C})| + |\mathcal{E}| + \sum_{i=0}^k |\mathcal{D}_i| \\ &= |f(\mathcal{B}) \cup f(\mathcal{C})| + |\mathcal{F}| \\ &\leq \binom{n - kr - 2}{r - 1} + \binom{n - kr - 2}{r - 2} = \binom{n - kr - 1}{r - 1} = |\mathcal{A}_1^*|. \end{aligned} \quad (2)$$

This completes the proof of the bound in Theorem 3.

We now need to show that if equality holds in (2) then $\mathcal{A} \simeq \mathcal{A}_1^*$. This follows in the same way as in the proof of Theorem 2 although the details are more involved. The only real difference is that for $k \geq 2$ the extremal family is always unique. The following two lemmas are obvious analogues of Lemmas 6 and 7.

Lemma 11 *If equality holds in (2) and the family $f(\mathcal{F})$ defined above is isomorphic to $\mathcal{B}_1^* = \{A \in [n - k - 1]_k^{(r-1)} : 1 \in A\}$ then either $\mathcal{A} \simeq \mathcal{A}_1^*$ or there exists $j \in [n]$ such that $\mathcal{A}_{j, j+k+1}^* = \{A \in [n]_k^{(r)} : j, j + k + 1 \in A\} \subset \mathcal{A}$.*

Lemma 12 *If equality holds in (2) and $A_{1,k+2}^* = \{A \in [n]_k^{(r)} : 1, k+2 \in A\} \subset \mathcal{A}$ then every set in $\mathcal{A}$ meets $\{1, k+2\}$. Moreover if $A = \{1, a_2, \dots, a_r\} \in [n]_k^{(r)}$ satisfies $A \cap \{1, k+2\} = \{1\}$ then exactly one of the sets A and $g(A) = \{k+2, a_2+k, \dots, a_r+k\}$ belongs to $\mathcal{A}$.*

Now suppose equality holds in (2). If $n = (k+1)r$ then clearly $\mathcal{A} \simeq \mathcal{A}_1^*$ so we may suppose that $n \geq (k+1)r + 1$. We will prove that $\mathcal{A} \simeq \mathcal{A}_1^*$ by induction on r . Clearly the result holds for $r = 1$ or 2 so suppose $r \geq 3$. Since equality holds in (2) we must have $|f(\mathcal{F})| = \binom{n-kr-2}{r-2}$ and so our inductive hypothesis for r implies that $f(\mathcal{F}) \simeq \mathcal{B}_1^* = \{A \in [n-k-1]_k^{(r-1)} : 1 \in A\}$. If $\mathcal{A} \not\simeq \mathcal{A}_1^*$ then Lemma 11 implies that there exists $j \in [n]$ such that $\mathcal{A}_{j,j+k+1}^* = \{A \in [n]_k^{(r)} : j, j+k+1 \in A\} \subset \mathcal{A}$. Without loss of generality we may suppose that $j = 1$.

By the first part of Lemma 12 we know that every set in $\mathcal{A}$ meets $\{1, k+2\}$. Hence we may write $\mathcal{A} = \mathcal{B}_1 \cup \mathcal{B}_{k+2} \cup \mathcal{B}_{1,k+2}$, where $\mathcal{B}_1 = \{A \in \mathcal{A} : 1 \in A, k+2 \notin A\}$, $\mathcal{B}_{k+2} = \{A \in \mathcal{A} : 1 \notin A, k+2 \in A\}$ and $\mathcal{B}_{1,k+2} = \{A \in \mathcal{A} : 1, k+2 \in A\}$. For uniqueness to hold we need to show that one of the families $\mathcal{B}_1$ or $\mathcal{B}_{k+2}$ must be empty. We will prove that if both families are non-empty then $\mathcal{A}$ contains two disjoint sets, a contradiction.

Since we are supposing that equality holds in (2), Lemma 12 tells us that if $A = \{1, a_2, \dots, a_r\} \in [n]_k^{(r)}$ and $A \cap \{1, k+2\} = \{1\}$ then exactly one of the sets A and $g(A) = \{k+2, a_2+k, \dots, a_r+k\}$ belong to $\mathcal{A}$. We now wish to show that if $A \in \mathcal{B}_1$ and $B \in [n]_k^{(r)}$ is obtained from A by shifting one vertex of $A \setminus \{1\}$ one place clockwise then $B \in \mathcal{B}_1$. Suppose not, then by Lemma 12 we have $g(B) \in \mathcal{B}_{k+2}$. But $g(B) \cap A = \emptyset$, contradicting the fact that $\mathcal{A}$ is intersecting. Similarly if $A \in \mathcal{B}_{k+2}$ and $B \in [n]_k^{(r)}$ is obtained from A by shifting one vertex of $A \setminus \{k+2\}$ one place anti-clockwise then $B \in \mathcal{B}_{k+2}$.

Now suppose, for a contradiction, that both $\mathcal{B}_1$ and $\mathcal{B}_{k+2}$ are non-empty. Then, by shifting points one by one, we see that $A = \{k+2, 2k+3, \dots, (k+1)r+1\}$ and $B = \{1, n-(k+1)(r-1)+1, \dots, n-2(k+1)+1, n-k\}$ both belong to $\mathcal{A}$. If $n \not\equiv 0 \pmod{k+1}$ then these are disjoint and we have a contradiction. So we may suppose that n is a multiple of $(k+1)$ and hence $n \geq (k+1)(r+1) \geq (k+1)r+2$. Then either there is a set in $\mathcal{B}_1$ not containing $n-k$ and so $C = \{1, n-(r-1)(k+1), \dots, n-2(k+1), n-(k+1)\} \in \mathcal{A}$ or $C \notin \mathcal{A}$. If $C \in \mathcal{A}$ then we have a contradiction since $A \cap C = \emptyset$. Otherwise Lemma 12 implies that $g(C) = \{k+2, n-(r-2)(k+1)-1, n-(r-3)(k+1)-1, \dots, n-1\} \in \mathcal{A}$. (Note that $C, g(C) \in [n]_k^{(r)}$ follows from $n \geq (k+1)r+2$.) But since $k \geq 2$ we have $1 \not\equiv -1 \pmod{k+1}$ and so $B \cap g(C) = \emptyset$. This contradiction completes

the proof of Theorem 3. $\square$

We now prove the lemmas. The definitions of $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{H}$ and f are as in the proof of Theorem 3.

Proof of Lemma 8. Let $A \in \mathcal{A}_1^* = \{A \in [n]_k^{(r)} : 1 \in A\}$, then A is described uniquely by the gaps $a_1, a_2, \dots, a_r \geq k$ such that

$$A = \{1, 2 + a_1, 3 + a_1 + a_2, \dots, r + a_1 + \dots + a_{r-1}\},$$

and $\sum_{i=1}^r a_i = n - r$. So $|\mathcal{A}_1^*|$ is equal to the number of ways of choosing integers $b_1, b_2, \dots, b_r \geq 0$ with $\sum_{i=1}^r b_i = n - (k+1)r$ which is simply

$$\binom{n - (k+1)r + r - 1}{r - 1} = \binom{n - kr - 1}{r - 1}.$$

$\square$

Proof of Lemma 9. Let $A, B \in [n]_k^{(r)}$ and suppose $f(A) = f(B)$ but $A \neq B$. If $a \geq 2$ then $a \in f^j(A) \iff a + j \in A$. Hence

$$A \cap \{j + 2, \dots, n\} = B \cap \{j + 2, \dots, n\}.$$

So if $A \neq B$ then there exist $c, d \in \{1, \dots, j + 1\}$ such that $c \in A$ and $d \in B$ and $c \neq d$. But $j \leq k$ implies that $A \cap \{1, \dots, j + 1\} = \{c\}$ and $B \cap \{1, \dots, j + 1\} = \{d\}$ or vice-versa and so $A \Delta B = \{c, d\}$ as required. $\square$

Proof of Lemma 10(a). Let $G \in f^{k-1}(\mathcal{E})$. Then there exists $E \in \mathcal{E}$ such that $f^{k-1}(E) = G$. So there exist $B \in \mathcal{B}$ and $C \in \mathcal{C}$ such that $f(B) = f(C) = E$. Now Lemma 9 with $j = 1$ implies that $B \Delta C = \{1, 2\}$. Then $1 \in C$ implies that $C \cap \{n - k + 1, \dots, n\} = \emptyset$. Hence $E \cap \{n - k, \dots, n\} = \emptyset$. So $G \cap \{n - 2k + 1, \dots, n\} = \emptyset$. Also $2 \in B \implies E \cap \{2, \dots, k + 1\} = \emptyset$ and we have

$$G \cap (\{n - 2k + 1, \dots, n - k\} \cup \{1, 2\}) = \{1\}. \quad (3)$$

Hence $G \setminus \{1\} \in [n - k]_k^{(r-1)}$.

Now suppose $G \in f^k(\mathcal{D}_0)$. Then there exists $D \in \mathcal{D}_0$ such that $G = f^k(D)$ and $1, k + 2 \in D$. Hence $1 \in G$ and

$$G \cap (\{n - 2k + 1, \dots, n - k\} \cup \{1, \dots, k + 2\}) = \{1, 2\} \quad (4)$$

So $G \setminus \{1\} \in [n - k]_k^{(r-1)}$.

Finally suppose that $1 \leq i \leq k$ and $G \in f^k(\mathcal{D}_i)$. Then there is $D \in \mathcal{D}_i$ such that $G = f^k(D)$ and $n+1-i, k+2-i \in D$. Hence $1 = f^k(k+2-i) \in G$ and

$$G \cap (\{n - (i+2k) + 1, \dots, n - k\} \cup \{1, \dots, k+2-i\}) = \{n+1-(i+k), 1\} \quad (5)$$

So again $G \setminus \{1\} \in [n-k]_k^{(r-1)}$. Hence $\mathcal{F} \subseteq [n-k]_k^{(r-1)}$. $\square$

Proof of Lemma 10(b). The fact that these families of sets are pairwise disjoint follows from considering how sets from these families meet the set $\{n-2k+1, \dots, n-k\} \cup \{2\}$.

First suppose $G \in f^{k-1}(\mathcal{E}) - \{1\}$. Using (3) we have

$$G \cap (\{n-2k+1, \dots, n-k\} \cup \{2\}) = \emptyset.$$

Now suppose $G \in f^k(\mathcal{D}_0) - \{1\}$. Using (4) we have

$$G \cap (\{n-2k+1, \dots, n-k\} \cup \{2\}) = \{2\}.$$

Finally suppose $G \in f^k(\mathcal{D}_i) - \{1\}$ with $1 \leq i \leq k$. Using (5) we have

$$G \cap (\{n-2k+1, \dots, n-k\} \cup \{2\}) = \{n+1-(i+k)\}.$$

Hence these families are pairwise disjoint. $\square$

Proof of Lemma 10(c). We wish to show that

$$\mathcal{F} = (f^{k-1}(\mathcal{E}) - \{1\}) \cup \bigcup_{i=0}^k (f^k(\mathcal{D}_i) - \{1\}),$$

is intersecting.

We will consider four cases. First suppose that $A, B \in f^k(\mathcal{D}_i) - \{1\}$. So there exist $D, E \in \mathcal{D}_i$ such that $f^k(D) \setminus \{1\} = A$ and $f^k(E) \setminus \{1\} = B$. If $i = 0$ then $k+2 \in D \cap E$ so $2 \in A \cap B$. Otherwise $1 \leq i \leq k$ and $n+1-i \in D \cap E$ so $n+1-(i+k) \in A \cap B$.

Next suppose that $A, B \in f^{k-1}(\mathcal{E}) - \{1\}$ then there exist $E, F \in \mathcal{E}$ such that $f^{k-1}(E) \setminus \{1\} = A$ and $f^{k-1}(F) \setminus \{1\} = B$. Now $E, F \in \mathcal{E}$ implies that there exist $B_1 \in \mathcal{B}$ and $C_1 \in \mathcal{C}$ such that $f(B_1) = E$ and $f(C_1) = F$. Then $B_1, C_1 \in \mathcal{A} \implies B_1 \cap C_1 \neq \emptyset$ and Lemma 9 implies that $2 \in B_1, 1 \in C_1$. So there exists $j \geq k+3$ such that $j \in B_1 \cap C_1$. Hence $3 \leq f^k(j) \in A \cap B$.

For the next case suppose $0 \leq i < j \leq k$, $A \in f^k(\mathcal{D}_i) - \{1\}$ and $B \in f^k(\mathcal{D}_j) - \{1\}$. Then there exist $C \in \mathcal{D}_i$ and $D \in \mathcal{D}_j$ such that $f^k(C) \setminus \{1\} = A$ and

$f^k(D) \setminus \{1\} = B$. Now $j \geq 1 \implies D \cap \{1, \dots, k+2\} = \{k+2-j\}$ and $k+2-j \notin C$. But $C, D \in \mathcal{A} \implies C \cap D \neq \emptyset$. Hence there exists $l \geq k+3$ such that $l \in C \cap D$ and so $3 \leq f^k(l) \in A \cap B$.

For the last case suppose that $A \in f^{k-1}(\mathcal{E}) - \{1\}$ and $B \in f^k(\mathcal{D}_i) - \{1\}$. Then there exist $D \in \mathcal{D}_i$ and $E \in \mathcal{E}$ such that $f^k(D) \setminus \{1\} = B$ and $f^{k-1}(E) \setminus \{1\} = A$. Furthermore there exist $B_1 \in \mathcal{B}$ and $C_1 \in \mathcal{C}$ such that $f(B_1) = f(C_1) = E$. Lemma 9 implies that $1 \in C_1$ and $2 \in B_1$. So $B_1 \cap \{1, \dots, k+2\} = \{2\}$ and $C_1 \cap \{1, \dots, k+1\} = \{1\}$. Also $D \in \mathcal{D}_i$ implies that

$$D \cap \{1, 2, \dots, k+2\} = \begin{cases} \{1, k+2\}, & i = 0 \\ \{k+2-i\}, & 1 \leq i \leq k. \end{cases}$$

Now $B_1, C_1, D \in \mathcal{A}$ implies that $B_1 \cap D$ and $C_1 \cap D$ are both nonempty. So if $i = 0$ then there exists $j \geq k+3$ such that $j \in B_1 \cap D$ and hence $3 \leq f^k(j) \in A \cap B$. Otherwise $1 \leq i \leq k$ and there exists $j \geq k+3$ such that $j \in C_1 \cap D$ and hence $3 \leq f^k(j) \in A \cap B$.

Hence $\mathcal{F}$ is intersecting. $\square$

Proof of Lemma 10(d). We need to prove that $f(\mathcal{F}) \subseteq [n-k-1]_k^{(r-1)}$. We note first that by part (a) of this lemma we have $\mathcal{F} \subseteq [n-k]_k^{(r-1)}$. Let $F \in \mathcal{F}$ and consider $f(F)$. Clearly $f(F) \in [n-k-1]^{(r-1)}$. We simply need to check that $f(F)$ does not contain a gap of size exactly $k-1$ (as a subset of $[n-k-1]$). This will follow if we show that F does not contain a gap of size exactly k (as a subset of $[n-k]$) around 1. More precisely we need to check that F does not contain any of the following pairs of points:

$$(n-2k+1, 2), \dots, (n-k, k+1), (1, k+2).$$

Since $1 \notin F$ we know that the last pair in this list cannot belong to F . Also using (3) and (4) we know that if $F \in f^{k-1}(\mathcal{E}) - \{1\}$ or $F \in f^k(\mathcal{D}_0) - \{1\}$ then

$$F \cap \{n-2k+1, \dots, n-k\} = \emptyset.$$

Finally using (5) we know that if $F \in f^k(\mathcal{D}_i) - \{1\}$, with $1 \leq i \leq k$, then

$$F \cap (\{n-2k+1, \dots, n-k\} \cup \{1, \dots, k+2-i\}) = \{n+1-(i+k)\}.$$

Hence if $n-2k+j \in F \in f^k(\mathcal{D}_i) - \{1\}$, with $1 \leq j \leq k$, then $j = k+1-i$ and so $j+1 = k+2-i \notin F$. Hence F cannot contain any of the pairs of points in the above list (since they are all of the form $(n-2k+j, j+1)$). So $f(\mathcal{F}) \subseteq [n-k-1]_k^{(r-1)}$ as required. $\square$

Proof of Lemma 11. Suppose that $f(\mathcal{F}) \simeq \mathcal{B}_1^* = \{A \in [n-k-1]_k^{(r-1)} : 1 \in A\}$. We need to show that either $\mathcal{A} \simeq \mathcal{A}_1^*$ or there exists $j \in [n]$ such that $\mathcal{A}_{j,j+k+1}^* = \{A \in [n]_k^{(r)} : j, j+k+1 \in A\} \subset \mathcal{A}$.

Since $f(\mathcal{F}) \simeq \mathcal{B}_1^*$ there exists $i \in [n-k-1]$ such that $f(\mathcal{F}) = \mathcal{B}_i^* = \{A \in [n-k-1]_k^{(r-1)} : i \in A\}$. If $i = 1$ then every set in $\mathcal{F}$ contains 2 and so $\mathcal{F} = f^k(\mathcal{D}_0) - \{1\}$. Since $\mathcal{D}_0 \subseteq \mathcal{A}_{1,k+2}^* = \{A \in [n]_k^{(r)} : 1, k+2 \in A\}$ and this last family has size $\binom{n-kr-2}{r-2} = |f(\mathcal{F})| = |\mathcal{D}_0|$ then $\mathcal{D}_0 = \mathcal{A}_{1,k+2}^*$. Similarly if $i \in \{n-2k, \dots, n-k-1\}$ then $\mathcal{A}_{j,j+k+1}^* \subset \mathcal{A}$, where $j = i+k+1$.

So suppose now that $i \in \{2, \dots, n-2k-1\}$. We will first show that $\mathcal{A}_{1,i+k+1}^* \cup \mathcal{A}_{2,i+k+1}^* \subseteq \mathcal{A}$. Suppose $A = \{1, a_2, \dots, a_r\} \in \mathcal{A}_{1,i+k+1}^*$ but $A \notin \mathcal{A}$. Then since $f(\mathcal{F}) = \mathcal{B}_i^*$ it must be the case that $B = \{a_2 - (k+1), \dots, a_r - (k+1)\} \in f(\mathcal{F})$. Hence $C = (A \setminus \{1\}) \cup \{j\} \in \mathcal{A}$ for some $j \in \{2, \dots, k+1\}$ and then $1 \in A$ implies that $C \cap \{n-k+1, \dots, n\} = \emptyset$. Hence $C \notin \bigcup_{i=0}^k \mathcal{D}_i$. This implies that $B \in f(f^{k-1}(\mathcal{E}) - \{1\})$ and so $A \in \mathcal{A}$, a contradiction. Hence $\mathcal{A}_{1,i+k+1}^* \subset \mathcal{A}$. Similarly $\mathcal{A}_{2,i+k+1}^* \subset \mathcal{A}$.

Define $\mathcal{C}_j^* = \{A \in [n]_k^{(r)} : i+k+1, j+1, n-k+j \in A\}$. So $\mathcal{D}_j \subseteq \mathcal{C}_{k+1-j}^*$. We claim that if $1 \leq j \leq k+1$ then $\mathcal{C}_j^* \subset \mathcal{A}$. Fix j and let $C \in \mathcal{C}_j^*$. Consider $B = f(f^k(C) \setminus \{1\}) \in \mathcal{B}_i^*$. Then $n-2k+j \in B$ implies that $C \in \mathcal{D}_{j-(k+1)} \subseteq \mathcal{A}$.

To summarise, we now know that

$$\mathcal{A}_{1,i+k+1}^* \cup \mathcal{A}_{2,i+k+1}^* \cup \bigcup_{j=1}^{k+1} \mathcal{C}_j^* \subseteq \mathcal{A}. \quad (6)$$

We can now show that every set in $\mathcal{A}$ must contain $i+k+1$ and so $\mathcal{A} \simeq \mathcal{A}_1^*$. Suppose not, then there exists a set $A = \{a_1, \dots, a_r\} \in \mathcal{A}$ such that $i+k+1 \notin A$. We will construct a set $B \in \mathcal{A}$ such that $A \cap B = \emptyset$.

Suppose $a_j < i+k+1 < a_{j+1}$. We consider first the case that $a_j = a_r$ and $a_{j+1} = a_1$. If $1 \leq i+k+1 < a_1$ then $i+k+1 \geq k+3$. So let $B = \{1, i+k+1, a_2-1, \dots, a_{r-1}-1\} \in \mathcal{A}_{1,i+k+1}^* \subset \mathcal{A}$. Then $A \cap B = \emptyset$. Otherwise we have $i+k+1 \leq n$ and hence $i+k+1 \leq n-k$. So let $B = \{2, a_2+1, \dots, a_{r-1}+1, i+k+1\}$ or $B = \{1, a_2+1, \dots, a_{r-1}+1, i+k+1\}$ depending on whether or not $a_1 = 1$. In either case $B \in \mathcal{A}_{1,i+k+1}^* \cup \mathcal{A}_{2,i+k+1}^* \subset \mathcal{A}$ and $A \cap B = \emptyset$.

We turn to the general case, i.e. $a_j < i+k+1 < a_{j+1}$ for $1 \leq j \leq r-1$. Let the gap between a_r and a_1 be of size $\alpha \geq k$, and suppose $a_j + \beta = a_{j+1} - \gamma = i+k+1$. We must consider two sub-cases. First suppose that $\max\{\beta, \gamma\} \leq k$. Let $C = \{a_1 + \beta, \dots, a_{j-1} + \beta, i+k+1, a_{j+2} - \gamma, \dots, a_r - \gamma\}$. So $i+k+1 \in C$ and $A \cap C = \emptyset$. The gap between the first and last elements of C is $\alpha + \beta + \gamma \geq 2k+1$.

If $a_r - \gamma + k + 1 \geq 1$ let $B = C \cup \{a_r - \gamma + k + 1\}$. Otherwise if $a_1 + \beta - (k + 1) \leq 2$ let $B = C \cup \{a_1 + \beta - (k + 1)\}$. If neither holds then let $B = C \cup \{1\}$. In each case $A \cap B = \emptyset$ and (6) implies that $B \in \mathcal{A}$ as required.

For the second sub-case we suppose $\max\{\beta, \gamma\} \geq k + 1$. Without loss of generality assume $\beta \geq k + 1$. Let $\delta = \max\{k + 1 - \gamma, 1\}$. We now define $C = \{a_1 - 1, \dots, a_j - 1, i + k + 1, a_{j+1} + \delta, \dots, a_{r-1} + \delta\}$. Now either $a_1 = 1$ in which case let $B = (C \setminus \{a_2 - 1\}) \cup \{k + 1\}$ or $a_1 \geq 2$. If $a_1 = 2$ or $a_1 = 3$ then let $B = C \in \mathcal{A}_{1,i+k+1}^* \cup \mathcal{A}_{2,i+k+1}^* \subseteq \mathcal{A}$ and $A \cap B = \emptyset$. So suppose $a_1 \geq 4$. In this case we construct B from C by shifting the last element of C clockwise and/or the first element of C anticlockwise until we obtain a set in $\cup_{j=1}^{k+1} \mathcal{C}_j^*$ that is disjoint from A . Again (6) implies that $B \in \mathcal{A}$ as required.

Hence every set in $\mathcal{A}$ contains $i + k + 1$ and so $\mathcal{A} \simeq \mathcal{A}_1^*$. $\square$

Proof of Lemma 12. Suppose equality holds in (2) and $\mathcal{A}_{1,k+2}^* \subseteq \mathcal{A}$. We show first that every set in $\mathcal{A}$ meets $\{1, k + 2\}$. Suppose not, then there is a set $A = \{a_1, \dots, a_r\} \in \mathcal{A}$ such that $A \cap \{1, k + 2\} = \emptyset$. Then $a_3 \geq 2k + 4$ so $B = \{1, k + 2, a_3 - 1, \dots, a_r - 1\} \in \mathcal{A}_{1,k+2}^* \subset \mathcal{A}$. But $A \cap B = \emptyset$ contradicting the fact that $\mathcal{A}$ is intersecting. Hence every set in $\mathcal{A}$ meets $\{1, k + 2\}$.

Let $\mathcal{B}_1 = \{A \in \mathcal{A} : 1 \in A, k + 2 \notin A\}$ and $\mathcal{B}_{k+2} = \{A \in \mathcal{A} : 1 \notin A, k + 2 \in A\}$. Then for equality to hold in (2) we must have $|\mathcal{B}_1 \cup \mathcal{B}_{k+2}| = \binom{n - kr - 2}{r - 1}$. Let $\mathcal{C}_1 = \{C \in [n]_k^{(r)} : 1 \in C, k + 2 \notin C\}$ and $\mathcal{C}_{k+2} = \{C \in [n]_k^{(r)} : 1 \notin C, k + 2 \in C\}$. Then for $A = \{1, a_2, \dots, a_r\} \in \mathcal{C}_1$ define $g(A) = \{k + 2, a_2 + k, \dots, a_r + k\}$. Clearly $A \cap g(A) = \emptyset$ and g is a bijection from $\mathcal{C}_1$ to $\mathcal{C}_{k+2}$. Then since $|\mathcal{C}_1| = |\mathcal{C}_{k+2}| = \binom{n - kr - 2}{r - 1}$ it must be the case that for any $A \in \mathcal{C}_1$ exactly one of the sets A and $g(A)$ belongs to $\mathcal{A}$. $\square$

4 The Erdős-Ko-Rado theorem for weighted k -separated sets

We conclude this paper with a short result for suitably weighted k -separated sets. Unlike our other results this follows simply from the original Erdős-Ko-Rado theorem (Theorem 1).

For $A = \{a_1, \dots, a_r\} \in [n]_k^{(r)}$ we define the *weight* of A to be

$$w(A) = \prod_{i=1}^r \binom{a_{i+1} - a_i - 1}{k},$$

where $a_{r+1} = a_1 + n$. So the weight of a k -separated set $A \in [n]_k^{(r)}$ is simply the number of different ways A may be extended to form a set $B \in [n]^{((k+1)r)}$ by inserting exactly k new elements into each gap in A . We then define the *weight* of a family of sets $\mathcal{A} \subseteq [n]_k^{(r)}$ to be

$$w(\mathcal{A}) = \sum_{A \in \mathcal{A}} w(A).$$

The following result says that an analogue of the Erdős-Ko-Rado theorem holds for weighted k -separated sets.

Theorem 4 *Let $n \geq 2(k+1)r$. If $\mathcal{A} \subseteq [n]_k^{(r)}$ is intersecting then $w(\mathcal{A}) \leq w(\mathcal{A}_1^*)$, where $\mathcal{A}_1^* = \{A \in [n]_k^{(r)} : 1 \in A\}$.*

Proof. Consider the bipartite graph $G = (V \cup W, E)$ with vertex classes $V = [n]_k^{(r)}$ and $W = [n]^{((k+1)r)}$. We define E as follows: let $A \in V$ be adjacent to $B \in W$ if we can construct B from A by inserting exactly k elements into each gap in A .

Let $\mathcal{A} \subseteq [n]_k^{(r)}$ be intersecting then

$$\Gamma(\mathcal{A}) = \{B \in [n]^{((k+1)r)} : \exists A \in \mathcal{A} \text{ such that } (A, B) \in E\}$$

is also intersecting. Then, since $\Gamma(\mathcal{A})$ is an intersecting family of $(k+1)r$ -sets from $[n]^{((k+1)r)}$ and $n \geq 2(k+1)r$, Theorem 1 implies that

$$|\Gamma(\mathcal{A})| \leq \binom{n-1}{(k+1)r-1}.$$

For distinct $A_1, A_2 \in \mathcal{A}$ we have $\Gamma(A_1) \cap \Gamma(A_2) = \emptyset$. To see this, suppose we had $B \in \Gamma(A_1) \cap \Gamma(A_2)$ with $B = \{b_1, \dots, b_{(k+1)r}\}$. Without loss of generality we may suppose that $A_1 = \{b_1, b_{k+2}, \dots, b_{(k+1)r-k}\}$ and $A_2 = \{b_i, b_{k+i+1}, \dots, b_{(k+1)(r-1)+i}\}$, for some $2 \leq i \leq k+1$. Hence $A_1 \cap A_2 = \emptyset$. This contradicts the fact that $\mathcal{A}$ is intersecting.

Then, since $|\Gamma(A)| = w(A)$, we have

$$w(\mathcal{A}) = \sum_{A \in \mathcal{A}} w(A) = \sum_{A \in \mathcal{A}} |\Gamma(A)| = |\Gamma(\mathcal{A})| \leq \binom{n-1}{(k+1)r-1} = w(\mathcal{A}_1^*).$$

□

References

- [1] P. ERDŐS, C. KO and R. RADO, ‘Intersection theorems for systems of finite sets’, *Quart. J. Math. Oxford (2)* 12 (1961), 313-320.

- [2] A.J.W. HILTON and E.C. MILNER, 'Some intersection theorems for systems of finite sets', *Quart. J. Math. Oxford (2)* 18 (1967), 369-384.
- [3] F.C. HOLROYD and A. JOHNSON, 'Problem 25', *Problems from the Sixteenth British Combinatorial Conference* edited by P.J. Cameron (1997), <http://www.maths.qmw.ac.uk/~pjc/bcc/allprobs.pdf>.
- [4] F.C. HOLROYD, 'Problem 338 (BCC16.25), Erdős-Ko-Rado at the court of King Arthur', *Discrete Mathematics* 197/198 (1999), 812.
- [5] M. KNESER, 'Aufgabe 360', *Jber. Deutsch. Math.-Verein.* 58 (1955/56), 2. Abt. 27.
- [6] L. LOVÁSZ, 'Kneser's conjecture, chromatic number, and homotopy', *J. Combinatorial Theory (A)* 25 (1978), 319-324.
- [7] A. SCHRIJVER, 'Vertex-critical subgraphs of Kneser graphs', *Nieuw Arch. Wisk. (3)* 26 (1978), 454-461.